





Identification of Novel *Acinetobacter baumannii* Host Fatty Acid Stress Adaptation Strategies

Jih-Hang Jiang,^a Karl A. Hassan,^b Stephanie L. Begg,^{c,d} Thusitha W. T. Rupasinghe,^e Varsha Naidu,^f Victoria G. Pederick,^c Marjan Khorvash,^c Jonathan J. Whittall,^{c,g} James C. Paton,^c Ian T. Paulsen,^f  Christopher A. McDewitt,^d Anton Y. Peleg,^{a,h}  Bart A. Eijkelkamp^c

^aInfection and Immunity Program, Monash Biomedicine Discovery Institute and Department of Microbiology, Monash University, Clayton, Victoria, Australia

^bSchool of Environmental and Life Sciences, University of Newcastle, Callaghan, New South Wales, Australia

^cResearch Centre for Infectious Diseases, School of Biological Sciences, University of Adelaide, Adelaide, South Australia, Australia

^dDepartment of Microbiology and Immunology, The Peter Doherty Institute for Infection and Immunity, University of Melbourne, Melbourne, Victoria, Australia

^eMetabolomics Australia, School of BioSciences, University of Melbourne, Melbourne, Victoria, Australia

^fDepartment of Chemistry and Biomolecular Sciences, Macquarie University, Sydney, New South Wales, Australia

^gSchool of Pharmacy and Medical Sciences, Sansom Institute for Health Research, University of South Australia, Adelaide, South Australia, Australia

^hDepartment of Infectious Diseases, The Alfred Hospital and Central Clinical School, Monash University, Melbourne, Victoria, Australia

ABSTRACT Free fatty acids hold important immune-modulatory roles during infection. However, the host's long-chain polyunsaturated fatty acids, not commonly found in the membranes of bacterial pathogens, also have significant broad-spectrum antibacterial potential. Of these, the omega-6 fatty acid arachidonic acid (AA) and the omega-3 fatty acid docosahexaenoic acid (DHA) are highly abundant; hence, we investigated their effects on the multidrug-resistant human pathogen *Acinetobacter baumannii*. Our analyses reveal that AA and DHA incorporate into the *A. baumannii* bacterial membrane and impact bacterial fitness and membrane integrity, with DHA having a more pronounced effect. Through transcriptional profiling and mutant analyses, we show that the *A. baumannii* β -oxidation pathway plays a protective role against AA and DHA, by limiting their incorporation into the phospholipids of the bacterial membrane. Furthermore, our study identified a second bacterial membrane protection system mediated by the AdelJK efflux system, which modulates the lipid content of the membrane via direct efflux of lipids other than AA and DHA, thereby providing a novel function for this major efflux system in *A. baumannii*. This is the first study to examine the antimicrobial effects of host fatty acids on *A. baumannii* and highlights the potential of AA and DHA to protect against *A. baumannii* infections.

IMPORTANCE A shift in the Western diet since the industrial revolution has resulted in a dramatic increase in the consumption of omega-6 fatty acids, with a concurrent decrease in the consumption of omega-3 fatty acids. This decrease in omega-3 fatty acid consumption has been associated with significant disease burden, including increased susceptibility to infectious diseases. Here we provide evidence that DHA, an omega-3 fatty acid, has superior antimicrobial effects upon the highly drug-resistant pathogen *Acinetobacter baumannii*, thereby providing insights into one of the potential health benefits of omega-3 fatty acids. The identification and characterization of two novel bacterial membrane protective mechanisms against host fatty acids provide important insights into *A. baumannii* adaptation during disease. Furthermore, we describe a novel role for the major multidrug efflux system AdelJK in *A. baumannii* membrane maintenance and lipid transport. This core function, beyond drug efflux, increases the appeal of AdelJK as a therapeutic target.

KEYWORDS AdelJK, antimicrobial host lipids, RND efflux, β -oxidation, free fatty acids, lipidomics

Citation Jiang J-H, Hassan KA, Begg SL, Rupasinghe TWT, Naidu V, Pederick VG, Khorvash M, Whittall JJ, Paton JC, Paulsen IT, McDewitt CA, Peleg AY, Eijkelkamp BA. 2019. Identification of novel *Acinetobacter baumannii* host fatty acid stress adaptation strategies. mBio 10:e02056-18. <https://doi.org/10.1128/mBio.02056-18>.

Editor Robert A. Bonomo, Louis Stokes Veterans Affairs Medical Center

Copyright © 2019 Jiang et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Bart A. Eijkelkamp, bart.eijkelkamp@adelaide.edu.au.

Received 20 September 2018

Accepted 12 December 2018

Published 5 February 2019




FIG 1 (A) The effects of long-chain polyunsaturated fatty acids (LC-PUFAs) on *A. baumannii* growth were determined by growing *A. baumannii* AB5075_UW in Luria-Bertani (LB) broth without LC-PUFA stress or with arachidonic acid (AA) or docosahexaenoic acid (DHA). The AA or DHA stress was quantified by comparing the 50% effective concentration (EC_{50}), which was calculated using Prism 8 (GraphPad) from the optical density at 600 nm (OD_{600}) measurements taken every 30 min, as per previous work (20). The EC_{50} is the time (minutes) when cultures have reached 50% of the maximum cell density, thereby representing mid-log growth. (B) The cell-associated omega-6 fatty acids (AA, GLA, and total n-6) and omega-3 fatty acids (DHA, EPA, and total n-3) were examined by gas chromatography following growth of strain AB5075_UW in the presence of 125 μ M AA- or DHA-supplemented LB, respectively, using routine methods (19, 20). (C) The modification of exogenous fatty acids into phospholipids, phosphatidylethanolamine (PE), and phosphatidylglycerol (PG), was quantified in AB5075_UW cells using liquid chromatography-mass spectrometry (LC-MS) and underwent further species verification by tandem MS (MS/MS) following published protocols (19, 21). (D) The membrane permeability of AB5075_UW and *adeJ121::T26* cells was examined by exposing the cells to 5 μ M Sytox for 5 min, followed by extensive washing and analysis on a PHERAstar spectrophotometer at excitation 485/emission 520 (BMG Labtech). (E) The effects of 125 μ M AA or DHA on *A. baumannii* AB5075_UW and the *fadB* and *fadA* Tn insertion mutant derivative *fadB167::T26* and *fadA140::T26* strains (14), respectively. The growth delay (in minutes) was calculated by comparing the EC_{50} s. (F) The conversion of exogenous DHA into cell-associated PE or PG phospholipids in AB5075_UW or *fadB167::T26* cells as determined by LC-MS. For all panels, the results are the mean \pm standard error of the mean (SEM) from at least biological triplicates. Statistical analyses were performed using a one-way analysis of variance (ANOVA) (A, C, D, and E) or a Student's *t* test (B and F). ns, not significant; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$.

Omega-3 and omega-6 fatty acids have distinct anti-*Acinetobacter* potential.

Considering the major disease burden associated with an imbalance in the human omega-3 and -6 fatty acid status (1, 2) and the well-documented antibacterial activity of long-chain polyunsaturated fatty acids (LC-PUFAs) (3, 4), we examined the impact of omega-3 and -6 fatty acids on *Acinetobacter baumannii*. *A. baumannii* is a human pathogen that is of primary concern in the hospital environment, where its exceptional capacity for antimicrobial resistance allows it to cause significant morbidity and mortality in susceptible patients (5, 6). Here, we investigated the effect of arachidonic acid (AA; 20:4n-6) or docosahexaenoic acid (DHA; 22:6n-3) at 31.25 to 500 μ M upon the growth of *A. baumannii* strain AB5075_UW. These concentrations and the magnitude of variation are physiologically relevant, considering differences greater than 10-fold in serum LC-PUFAs between individuals can be seen, depending largely on dietary intake (7). We found that both LC-PUFAs induced growth perturbation, but this was more pronounced with DHA compared to AA exposure (Fig. 1A). Interestingly, preexposure to DHA did not affect the cell's subsequent growth dynamics in the presence of DHA (see Fig. S1A in the supplemental material). Examination of alternative LC-PUFA fatty

acids revealed that the antimicrobial potential of omega-3 and -6 fatty acids may be conserved because γ -linolenic acid (GLA; omega-6) and eicosapentaenoic acid (EPA; omega-3) have similar impacts on the growth of strain AB5075_UW as AA and DHA, respectively (Fig. S1B and C).

We next examined the relative accumulation of the exogenous fatty acids in *A. baumannii* upon *in vitro* treatment with either 125 μ M AA or 125 μ M DHA, representing the lowest concentration at which the omega-3 fatty acid shows a significantly greater antimicrobial potential compared to the omega-6 fatty acid (Fig. 1A; Fig S1B and C). This revealed that not only are the exogenous fatty acids incorporated, but *A. baumannii* partly degrades AA and DHA, generating GLA (18:3n-6) and EPA (20:5n-3), respectively (Fig. 1B). Examination of the two major phospholipid species phosphatidylethanolamine (PE) and phosphatidylglycerol (PG) suggested that the exogenous lipids were readily incorporated into phospholipids (28% and 20% of total phospholipids after AA and DHA treatment, respectively) (Fig. 1C). This strongly indicates that the bacterium expresses systems that actively interact with host fatty acids. Based on the level of host fatty acid incorporation into the bacterial membrane, we investigated the impact of AA or DHA on bacterial membrane integrity (Fig. 1D). Treatment with AA or DHA increased the membrane permeability, but this was most significant (3.1-fold) after exposure to DHA (Fig. 1D). Overall, despite AA accumulating in the bacterial cell at higher levels, the longer and more desaturated host fatty acid DHA exerted a greater effect on *A. baumannii* growth and membrane integrity.

Membrane phospholipids play a key role in the defense against antimicrobials, including host fatty acids (8). Hence, we examined the effect of AA or DHA treatment upon the abundance of the endogenous *A. baumannii* fatty acids. The results showed a highly specific depletion ($\geq 50\%$) of the monounsaturated fatty acids 16:1n-7 and 18:1n-9 (Fig. S1D). This could not be accounted for through the transcriptional dysregulation of the FASII pathway, as seen in Gram-positive species (9). However, the *fadAB* operon (ABUW_3573-3572) was significantly upregulated upon treatment with AA or DHA (Fig. S1E). The *fadAB* operon encodes critical components of the fatty acid β -oxidation pathway, indicating fatty acid degradation may provide protection against LC-PUFAs. Indeed, the fitness of *fadA*- and *fadB*-inactivated mutants was significantly compromised under AA and DHA stress (Fig. 1E). Interestingly, the accumulation of DHA or EPA was not affected by mutation of *fadB*, despite its hypersusceptibility to DHA (Fig. S1F). Instead, we observed that of the PE and PG species with exogenous fatty acid incorporated, PE(36:5) and PG(36:6), the levels were significantly higher in the *fadB* mutant strain upon treatment with DHA (Fig. 1F). This suggests that the β -oxidation pathway is likely to restrict conversion of DHA and its derivatives into phospholipids. Collectively, these data indicate that the β -oxidation pathway contributes to protection against LC-PUFAs, which may impact the success of *A. baumannii* as a human pathogen.

The AdelJK RND efflux system is involved in LC-PUFA resistance. The *A. baumannii* resistance nodulation cell division (RND) family of efflux systems has been associated with *in vivo* survival (10). Hence, we first studied their involvement in LC-PUFA stress resistance by examining the transcriptional responsiveness upon AA or DHA supplementation. Interestingly, AA and DHA induced the specific upregulation of *adeJ* (Fig. 2A). *adeJ* is part of the *A. baumannii* core genome and, in combination with *adel* and *adeK*, encodes the complete RND efflux system (AdelJK) that plays an important role in *A. baumannii* multidrug resistance (10, 11). In recent years, several studies have revealed the RND efflux systems of *Acinetobacter* species, including AdelJK, to play roles in virulence and virulence-associated phenotypes (11, 12). Although their exact mode of action remains unknown, it has been postulated to involve lipid homeostasis (13). We examined the effect of *adeJ* inactivation on susceptibility to AA and DHA in both *A. baumannii* strain AB5075_UW (14) and *Acinetobacter baylyi* strain ADP1 (15). Growth delays were observed in both *adeJ* mutants (Fig. 2B), confirming that AdelJK plays a role in protection against AA and DHA and that this function is conserved between at least two different *Acinetobacter* species. Expression of *adelJK* is repressed by AdeN (16);




FIG 2 (A) Transcription levels of genes that encode putative *A. baumannii* resistance nodulation cell division (RND) efflux pumps in AB5075_UW cells grown in LB media with or without AA or DHA were examined by quantitative reverse transcription-PCR (qRT-PCR), using the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene as a reference gene, and data were corrected against untreated cells as per previous analyses (19, 20). AB5075_UW cultures were grown to an optical density at 600 nm (OD_{600}) of 0.5 and were exposed to 125 μ M AA or DHA for 30 min prior to processing for RNA isolation. Oligonucleotide sequences can be found in Table S1 in the supplemental material. The results are the mean \pm SEM from at least biological triplicates. Statistical analyses were performed using a one-sample *t* test (**, $P < 0.01$; ***, $P < 0.001$). (B) The effect of 125 μ M AA or DHA on *adeJ*-inactivated *A. baumannii* AB5075_UW (*adeJ121::T26*) and *A. baylyi* ADP1 (Δ *adeJ*) mutants (14, 15). The growth delay (in minutes) was calculated by comparing the 50% effective concentration (EC_{50}) (20). The results are the mean \pm SEM from at least biological triplicates. Statistical analyses were performed using a one-way ANOVA (*, $P < 0.05$; ***, $P < 0.001$; ****, $P < 0.0001$). (C) Endogenous phosphatidylglycerol (PG) and cardiolipin (CL) species that were significantly affected by mutation of *adeJ* in strain AB5075_UW (*adeJ121::T26*). The results are the mean \pm SEM from at least biological triplicates. Statistical analyses were performed using a two-way ANOVA on all PG or CL species identified by LC-MS, which can be found in Table S2 in the supplemental material (****, $P < 0.001$; *****, $P < 0.0001$). (D) The membrane permeability of AB5075_UW and *adeJ121::T26* cells was examined by exposing the cells to 5 μ M Sytox for 5 min, followed by extensive washing and analysis on a PHERAstar spectrophotometer at excitation 485/emission 520 (BMG Labtech). The results are the mean \pm SEM from at least biological triplicates. Statistical analyses were performed using a Student's *t* test (*, $P < 0.05$). (E) The fatty acids in the media were analyzed by gas chromatography following growth of the AB5075_UW or *adeJ121::T26* strains. The baseline represents media in which no bacteria were grown. Samples were first centrifuged at low speed ($300 \times g$ for 5 min), with the supernatant then passed through a 0.4 μ m-pore filter and insoluble material subsequently removed by ultracentrifugation ($100,000 \times g$ for 1 h). The results are the mean \pm SEM from at least biological triplicates. Statistical analyses were performed using a Student's *t* test (*, $P < 0.05$). (F) The ability of AB5075_UW or *adeJ121::T26* cells to form biofilms was assessed after 24 h of static growth in a polystyrene 96-well plate. Cells were stained using 0.1% crystal violet, with the absorbance determined at 590 nm (21, 22). The results are the mean \pm SEM from at least biological triplicates. Statistical analyses were performed using a Student's *t* test (*, $P < 0.05$). (G) Surface motility of AB5075_UW or *adeJ121::T26* cells was examined by inoculating the center of a semisolid LB plate (0.25% agar) and incubating for 18 h (22). The aberrant motility phenotype was observed in all 3 independent experiments (replicates 1, 2, and 3).

hence, we analyzed the growth dynamics of an *adeN* mutation in *A. baumannii*. Consistent with previous reports (10), the growth rate of the *adeN* mutant is compromised compared to that of the parental strain (see Fig. S2A in the supplemental material). However, growth rates in the presence of DHA were similar between the *adeN* mutant and wild type (WT) (Fig. S2A), indicating that DHA-mediated *adeIJK* derepression provides *A. baumannii* with DHA resistance at a level similar to that when *adeN* has been inactivated. We then hypothesized that the *AdelJK* RND efflux system was responsible for the export of the exogenous fatty acid as a protection mechanism. Hence, we examined the accumulation of DHA in the *adeJ* mutant, but found that the accumulation of DHA and its derivative EPA, as well as the conversion into phospholipids, was significantly lower in the mutant compared to the parental strain (Fig. S2B and C). We then ascertained whether *AdelJK* may be involved in membrane modulation by removing endogenous fatty acids from the membrane to achieve lipid homeostasis, similar to the EmhABC RND efflux system of *Pseudomonas fluorescens* (17). Lipid

analyses demonstrated that the concentrations of two endogenous species of PG and two of cardiolipin were affected by *adeJ* mutation prior to treatment (Fig. 2C), with eight species in total being affected posttreatment with DHA (Fig. S2D). These findings implicate a role for AdeJ in membrane modulation. Indeed, even without treatment, AdeJ-mediated changes in the phospholipids affected the membrane integrity, as the permeability of the *adeJ* mutant was significantly greater than that of the WT (Fig. 2D). We then examined the fatty acids in the media following growth of the WT or *adeJ* mutant (Fig. 2E). Consistent with a role for AdeJ in lipid efflux, growth of the WT resulted in an increase in fatty acids in the media, whereas growth of the mutant strain resulted in fatty acid depletion (Fig. 2E). Hence, we speculate that the differences seen between the WT and *adeJ* mutant could most likely be ascribed to AdeJ-mediated export of fatty acids—a significant observation that indicates a novel role for AdeJ. We also found that mutation of *adeJ* leads to an increase in biofilm formation but a decrease in surface motility in *A. baumannii* (Fig. 2F and G). Thus, our data suggest that the exported lipids may function as a surfactant, which renders the biofilm unstable, but promotes surface migration. Here, we show for the first time, through its identification as a protective mechanism against LC-PUFAs, that the AdeJ RND efflux system is directly involved in the maintenance of lipid homeostasis in *A. baumannii*. Furthermore, through its proposed role as a lipid export system, this work has significantly advanced our understanding of the molecular basis behind AdeJ's far-reaching impacts on *A. baumannii* biofilm formation and *in vivo* fitness. Furthermore, similar to the lipid-mediated sequestration of daptomycin in *Staphylococcus aureus* (18, 19), the export of lipids may serve as a novel mechanism by which AdeJ provides protection against amphiphilic or hydrophobic antimicrobials.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/mBio.02056-18>.

FIG S1, TIF file, 0.4 MB.

FIG S2, TIF file, 0.4 MB.

TABLE S1, DOCX file, 0.1 MB.

TABLE S2, XLSX file, 0.1 MB.

ACKNOWLEDGMENTS

This work was supported by the National Health and Medical Research Council (Australia) through project grants 1080784 and 1122582 to C.A.M. and program grant 1071659 to J.C.P. The work was also funded by the Australian Research Council (ARC) Discovery project grant DP170102102 to J.C.P. and C.A.M. Work performed by K.A.H. and I.T.P. is supported by NHMRC project grant 1120298. B.A.E. is a University of Adelaide Beacon Research Fellow, S.L.B. is an NHMRC Doherty Fellow (1142695), C.A.M. is an ARC Future Fellow (FT170100006), and A.Y.P. is an NHMRC Practitioner Fellow (1117940).

REFERENCES

- Anes E, Jordao L. 2008. Trick-or-treat: dietary lipids and host resistance to infectious disease. *Mini Rev Med Chem* 8:1452–1458.
- Simopoulos AP. 2011. Evolutionary aspects of diet: the omega-6/omega-3 ratio and the brain. *Mol Neurobiol* 44:203–215. <https://doi.org/10.1007/s12035-010-8162-0>.
- Desbois AP, Smith VJ. 2010. Antibacterial free fatty acids: activities, mechanisms of action and biotechnological potential. *Appl Microbiol Biotechnol* 85:1629–1642. <https://doi.org/10.1007/s00253-009-2355-3>.
- Churchward CP, Alany RG, Snyder LAS. 2018. Alternative antimicrobials: the properties of fatty acids and monoglycerides. *Crit Rev Microbiol* 44:561–570. <https://doi.org/10.1080/1040841X.2018.1467875>.
- Harding CM, Hennon SW, Feldman MF. 2018. Uncovering the mechanisms of *Acinetobacter baumannii* virulence. *Nat Rev Microbiol* 16: 91–102. <https://doi.org/10.1038/nrmicro.2017.148>.
- Dijkshoorn L, Nemec A, Seifert H. 2007. An increasing threat in hospitals: multidrug-resistant *Acinetobacter baumannii*. *Nat Rev Microbiol* 5:939–951. <https://doi.org/10.1038/nrmicro1789>.
- Abdelmagid SA, Clarke SE, Nielsen DE, Badawi A, El-Sohehy A, Mutch DM, Ma DWL. 2015. Comprehensive profiling of plasma fatty acid concentrations in young healthy Canadian adults. *PLoS One* 10:e0116195. <https://doi.org/10.1371/journal.pone.0116195>.
- Eijkelkamp BA, Begg SL, Pederick VG, Trapetti C, Gregory MK, Whittall JJ, Paton JC, McDevitt CA. 2018. Arachidonic acid stress impacts pneumococcal fatty acid homeostasis. *Front Microbiol* 9:813. <https://doi.org/10.3389/fmicb.2018.00813>.
- Yao J, Rock CO. 2017. Exogenous fatty acid metabolism in bacteria. *Biochimie* 141:30–39. <https://doi.org/10.1016/j.biochi.2017.06.015>.
- Yoon E-J, Chabane YN, Goussard S, Snesrud E, Courvalin P, Dé E, Grillot-Courvalin C. 2015. Contribution of resistance-nodulation-cell division efflux systems to antibiotic resistance and biofilm formation

- in *Acinetobacter baumannii*. mBio 6:e00309-15. <https://doi.org/10.1128/mBio.00309-15>.
11. Yoon E-J, Balloy V, Fiette L, Chignard M, Courvalin P, Grillot-Courvalin C. 2016. Contribution of the Ade resistance-nodulation-cell division-type efflux pumps to fitness and pathogenesis of *Acinetobacter baumannii*. mBio 7:e00697-16. <https://doi.org/10.1128/mBio.00697-16>.
 12. Knight DB, Rudin SD, Bonomo RA, Rather PN. 2018. *Acinetobacter nosocomialis*: defining the role of efflux pumps in resistance to antimicrobial therapy, surface motility, and biofilm formation. Front Microbiol 9:1902. <https://doi.org/10.3389/fmicb.2018.01902>.
 13. Leus IV, Weeks JW, Bonifay V, Smith L, Richardson S, Zgurskaya HI. 2018. Substrate specificities and efflux efficiencies of RND efflux pumps of *Acinetobacter baumannii*. J Bacteriol 200:e00049-18. <https://doi.org/10.1128/JB.00049-18>.
 14. Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, Huse HK, Zurawski DV, Brittnacher MJ, Manoil C. 2015. Resources for genetic and genomic analysis of emerging pathogen *Acinetobacter baumannii*. J Bacteriol 197:2027–2035. <https://doi.org/10.1128/JB.00131-15>.
 15. Brzoska AJ, Hassan KA, de Leon EJ, Paulsen IT, Lewis PJ. 2013. Single-step selection of drug resistant *Acinetobacter baylyi* ADP1 mutants reveals a functional redundancy in the recruitment of multidrug efflux systems. PLoS One 8:e56090. <https://doi.org/10.1371/journal.pone.0056090>.
 16. Rosenfeld N, Bouchier C, Courvalin P, Périchon B. 2012. Expression of the resistance-nodulation-cell division pump AdeJJK in *Acinetobacter baumannii* is regulated by AdeN, a TetR-type regulator. Antimicrob Agents Chemother 56:2504–2510. <https://doi.org/10.1128/AAC.06422-11>.
 17. Adebosuyi AA, Foght JM. 2011. An alternative physiological role for the EmhABC efflux pump in *Pseudomonas fluorescens* cLP6a. BMC Microbiol 11:252. <https://doi.org/10.1186/1471-2180-11-252>.
 18. Sabnis A, Ledger EVK, Pader V, Edwards AM. 2018. Antibiotic interceptors: creating safe spaces for bacteria. PLoS Pathog 14:e1006924. <https://doi.org/10.1371/journal.ppat.1006924>.
 19. Pader V, Hakim S, Painter KL, Wigneshweraraj S, Clarke TB, Edwards AM. 2016. *Staphylococcus aureus* inactivates daptomycin by releasing membrane phospholipids. Nat Microbiol 2:16194. <https://doi.org/10.1038/nmicrobiol.2016.194>.
 20. Hassan KA, Pederick VG, Elbourne LDH, Paulsen IT, Paton JC, McDevitt CA, Eijkelkamp BA. 2017. Zinc stress induces copper depletion in *Acinetobacter baumannii*. BMC Microbiol 17:59. <https://doi.org/10.1186/s12866-017-0965-y>.
 21. Botté CY, Yamaryo-Botté Y, Rupasinghe TWT, Mullin KA, MacRae JI, Spurck TP, Kalanon M, Shears MJ, Coppel RL, Crellin PK, Maréchal E, McConville MJ, McFadden GI. 2013. Atypical lipid composition in the purified relict plastid (apicoplast) of malaria parasites. Proc Natl Acad Sci U S A 110:7506–7511. <https://doi.org/10.1073/pnas.1301251110>.
 22. Eijkelkamp BA, Stroehrer UH, Hassan KA, Papadimitriou MS, Paulsen IT, Brown MH. 2011. Adherence and motility characteristics of clinical *Acinetobacter baumannii* isolates. FEMS Microbiol Lett 323:44–51. <https://doi.org/10.1111/j.1574-6968.2011.02362.x>.