



## Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

# Mitochondrial genome of the Silvertip shark, *Carcharhinus albimarginatus*, from the British Indian Ocean Territory

Shaili Johri, Nicholas Dunn, Taylor K. Chapple, David Curnick, Vincent Savolainen, Elizabeth A. Dinsdale & Barbara A. Block

To cite this article: Shaili Johri, Nicholas Dunn, Taylor K. Chapple, David Curnick, Vincent Savolainen, Elizabeth A. Dinsdale & Barbara A. Block (2020) Mitochondrial genome of the Silvertip shark, *Carcharhinus albimarginatus*, from the British Indian Ocean Territory, Mitochondrial DNA Part B, 5:3, 2085-2086, DOI: [10.1080/23802359.2020.1765210](https://doi.org/10.1080/23802359.2020.1765210)

To link to this article: <https://doi.org/10.1080/23802359.2020.1765210>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 14 May 2020.



Submit your article to this journal [↗](#)



Article views: 387



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

## Mitochondrial genome of the Silvertip shark, *Carcharhinus albimarginatus*, from the British Indian Ocean Territory

Shaili Johri<sup>a,b</sup>, Nicholas Dunn<sup>c,d</sup>, Taylor K. Chapple<sup>e</sup>, David Curnick<sup>c</sup>, Vincent Savolainen<sup>d</sup>, Elizabeth A. Dinsdale<sup>b</sup> and Barbara A. Block<sup>a</sup>

<sup>a</sup>Hopkins Marine Station, Stanford University, Pacific Grove, CA, USA; <sup>b</sup>Department of Biology, San Diego State University, San Diego, CA, USA; <sup>c</sup>Zoological Society of London, Institute of Zoology, London, UK; <sup>d</sup>Department of Life Sciences, Imperial College London, Ascot, UK; <sup>e</sup>Coastal Oregon Marine Experiment Station, Oregon State University, Newport, OR, USA

### ABSTRACT

The Chagos archipelago in the British Indian Ocean Territory (BIOT) has been lacking in detailed genetic studies of its chondrichthyan populations. Chondrichthyes in Chagos continue to be endangered through illegal fishing operations, necessitating species distribution and abundance studies to facilitate urgent monitoring and conservation of the species. Here, we present a complete mitochondrial genome of the Silvertip Shark, *Carcharhinus albimarginatus* sampled in the Chagos archipelago. The mitochondrial genome of *C. albimarginatus* was 16,706 bp in length and consisted of 13 protein-coding genes, 22 tRNA genes, two rRNA genes, a replication origin and a D-loop region. GC content was at 38.7% and the control region was 1,065 bp in length. We expect that mitogenomes presented here will aid development of molecular assays for species distribution studies. Overall these studies will promote effective conservation of marine ecosystems in the BIOT.

### ARTICLE HISTORY

Received 27 March 2020  
Accepted 25 April 2020

### KEYWORDS

Chagos; MPA; sharks; ecosystems; conservation; MinION; molecular taxonomy



The Silvertip shark (*Carcharhinus albimarginatus*) is a large requiem shark species with fragmented distribution in the tropical Indian and Pacific Oceans, and is currently listed as 'Vulnerable' on the IUCN Red List (González-Medina and Pillans 2015). It is found throughout the British Indian Ocean Territory Marine Protected Area (BIOT MPA), but is frequently caught by illegal fisheries (Ferretti et al. 2018; Tickler et al. 2019), leading to severe population declines locally (Ferretti et al. 2018).

We sequenced the mitochondrial genome of *C. albimarginatus*. Samples were obtained from two individuals in the BIOT in March 2018 (Specimen 1: Latitude: 07°08'50.1600"S; Longitude: 072°08'19.5000"E and Specimen 2: Latitude: 07°08'17.6400"S; Longitude: 072°11'48.1800"E). These specimens were stored at the Hopkins Marine Station, Stanford University (Sample Accession #0200022322322) and Silwood Park, Imperial College London (Sample Accession #020002086005) in 70% ethanol. DNA for Specimen 1 was extracted from the fin clip of a juvenile male and sequenced on the MinION handheld sequencer following (Johri et al. 2019). Ninety-five Fast5 files obtained from 72 h of sequencing were converted to FASTQ files using the Guppy 3.3.1 basecaller (Oxford Nanopore Technologies, ONT) and a GPU interface for 12 h post sequencing. A total of 374, 396 sequence reads were obtained from the sequencing run, with a length range of 350 to 65,000 base pairs (bp). The reads were trimmed and mapped as described previously (Johri et al. 2019). The resulting contig consisted of 115 mapped reads which were checked for nucleotide assignment and alignment errors, and highest Q scores were used


to build consensus at ambiguous sites. The consensus sequence was annotated following (Johri et al. 2019; Johri et al. 2020). Second, genomic DNA was extracted from another juvenile male (specimen 2) using the Qiagen Blood & Tissue Kit and sequenced using an Illumina HiSeq platform. The mitochondrial genome sequence was then extracted bio-informatically yielding an almost identical sequence, which only differed from the sequence for Specimen 1 by 3 bp. These sequence differences include 3 single nucleotide polymorphisms which may be actual sequence differences or artifacts resulting from sequencing methods.

The mitochondrial genome of *C. albimarginatus* (GenBank: MT104516, MT093206) was 16,706 bp in length and consisted of 13 protein-coding genes (PCGs), 22 tRNA genes, two rRNA genes, a replication origin and a D-loop region. GC content was at 38.7%. All PCGs started with ATG and some PCGs ended with an incomplete stop codon. The control region was 1,065 bp in length.

To assess the phylogenetic position of *C. albimarginatus*, gene trees were constructed using mitogenomes from six families within the order Carcharhiniiformes and Hexanchiformes was used as outgroup. Phylogenies were determined in Bayesian inference frameworks (Huelsenbeck and Ronquist 2001; Edgar 2004; Ronquist et al. 2012) following (Johri et al. 2019; Johri et al. 2020). Bayesian trees were estimated using the GTR substitution model, gamma rate variations with 4 gamma categories, chain length 110,000, burn-in length 100,000 and subsampling frequency 200. *C. albimarginatus*

**CONTACT** Barbara A. Block  [bblock@stanford.edu](mailto:bblock@stanford.edu)  Hopkins Marine Station, Stanford University, 120 Ocean View Blvd, Pacific Grove, CA 93950, USA

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1.** Bayesian estimate of phylogenetic position of *Carcharhinus albimarginatus* within the order Carcharhiniformes computed with complete mitochondrial genomes. Members of the order Hexanchiformes served as the outgroup. Families are indicated by vertical lines and orders by square brackets. Numbers at nodes are posterior probabilities. GenBank Accession Numbers: *Notorynchus cepedianus* (AB560489.1); *Cephaloscyllium umbratile* (KT003686.1); *Scyliorhinus canicula* (Y16067.1); *Proscyllium habereri* (KU721838.1); *Pseudotriakis microdon* (AB560493.1); *Hemipristis elongata* (KU508621.1); *Hemigaleus microstoma* (KT003687.1); *Lamiopsis temminckii* (KT698048.1); *Glyphis fowlerae* (KT698049.1); *G. gangeticus* (KT698040.1); *Carcharhinus melanopterus* (KJ720818.1); *C. sorrah* (KF612341.1); *C. falciformis* (MK092088); *C. acronotus* (KF728380.1); *Carcharhinus albimarginatus* \_Ref (JQ518609.1); *C. albimarginatus* \_C (MT104516); *Eusphyra blochii* (KU892590.1); *Sphyrna lewini* (JX827259.1).

was nested within Carcharhinidae and most closely related to the *C. albimarginatus* reference (Figure 1).

Mitogenomes presented here will enable identification of interspecies genetic divergence through PCR amplification and accurate species identification using environmental DNA sampling. The current research findings will enable substantial conservation research and management of Chondrichthyes in the BIOT MPA and in other marine ecosystems.

## Acknowledgements

We thank the Stanford University, the Bertarelli Foundation, and the UK Natural Environment Research Council for funding, Robert Edwards and Adrian Cantu for computing support, Danny Coffey, Tim White, Nathan Truelove, Robert Schallert, and David Jacoby for collecting samples, Theodore Reimer for logistics and CD-genomics for assistance with Illumina sequencing.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Data availability

Data that support the findings of this study are openly available in Genbank with reference accession numbers MT104516 and MT093206 at DOI: <https://www.ncbi.nlm.nih.gov/nuccore/MT104516.1> and <https://www.ncbi.nlm.nih.gov/nuccore/MT093206.1>, respectively.

## Author contributions

SJ contributed to concept, sequencing, bioinformatics and wrote the manuscript with comments from VS, ND contributed to assembly and

bioinformatic analyses, TC and DC contributed to sample collection, ED provided laboratory support and BB contributed to the concept and logistical support for completion of the project. All authors reviewed and edited the manuscript.

## References

- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32(5):1792–1797.
- Ferretti F, Curnick D, Liu K, Romanov EV, Block BA. 2018. Shark baselines and the conservation role of remote coral reef ecosystems. *Sci Adv.* 4(3):eaq0333.
- González-Medina E, Pillans QR (CSIRO, Q. Mario Espinoza (James Cook University), and C./I. S. S. G. Nicholas Dulvy (Simon Fraser University). 2015. IUCN Red List of Threatened Species: Silvertip Shark. IUCN Red List of Threatened Species. <https://www.iucnredlist.org/en>.
- Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics.* 17(8):754–755.
- Johri S, Fellows SR, Solanki J, Busch A, Livingston I, Mora MF, Tiwari A, Cantu VA, Goodman A, Morris MM, et al. 2020. Mitochondrial genome to aid species delimitation and effective conservation of the Sharpnose Guitarfish (*Glaucostegus granulatus*). *Meta Gene.* 24:100648.
- Johri S, Solanki J, Cantu VA, Fellows SR, Edwards RA, Moreno I, Vyas A, Dinsdale EA. 2019. Genome skimming' with the MinION hand-held sequencer identifies CITES-listed shark species in India's exports market. *Sci Rep.* 9(1):4476.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP, et al. 2012. MrBayes 3.2: efficient bayesian phylogenetic inference and model choice across a large model space. *Syst Biol.* 61(3):539–542.
- Tickler DM, Carlisle AB, Chapple TK, Curnick DJ, Dale JJ, Schallert RJ, Block BA. 2019. Potential detection of illegal fishing by passive acoustic telemetry. *Anim Biotelemetry.* 7(1):1.