

# A literature review on plant-based foods and dietary quality in knee osteoarthritis

Zhaoli Dai<sup>1,2</sup>

## Abstract

This literature review summarizes the role of plant-based foods and diet quality in osteoarthritis, particularly knee osteoarthritis, in observational studies and clinical trials published during 2015–2020. The included studies have suggested favorable results on reducing the prevalence, pain, and cartilage changes related to osteoarthritis and inflammatory and oxidation markers such as interleukin-1, interleukin-6, tumor necrosis factor, and lipid peroxidation. Due to the lack of large longitudinal cohorts to study whole foods or diets concerning knee osteoarthritis, findings from the cross-sectional studies or clinical trials require further validation, particularly in well-designed clinical trials and a more extended follow-up period. Potential mechanisms on the role of plant-based foods in body weight, inflammation, and microbiome were explored to explain their protective associations with osteoarthritis. However, most evidence examining the relationship between the microbiome and osteoarthritis joint pain is conducted in preclinical animal studies, and few observational studies show a positive association between *Streptococcus* species and local joint inflammation in the knee. Given the close links of plant-based foods on obesity, inflammation, and microbiome, data on the role of whole foods or diets in the change in knee osteoarthritis pain through the lens of microbial composition can provide more certainty regarding the utilization of microbiome as a potential therapeutic target.

**Keywords:** Plant-based foods, dietary patterns, knee osteoarthritis, osteoarthritis

## Introduction

Knee osteoarthritis (OA), affecting 330 million people globally, is a leading cause of functional limitation and disability.<sup>1,2</sup> With global aging and nearly 40% of the world population being overweight and obese,<sup>3</sup> OA is a common age-related chronic condition that contributes to the global burden of disease.<sup>4</sup> Moreover, a well-established risk factor, overweight/obesity, significantly increases the risk of the onset and progression of OA.<sup>5</sup> This increased risk is partly due to bodyweight affecting the joints through mechanical loading. Furthermore, excessive adipose tissue also increases oxidative stress and pro-inflammatory markers such as tumor necrosis factor (TNF)  $\alpha$  and interleukin (IL)-6 locally in the joints and systemically in the blood.<sup>6,7</sup>

People with knee OA often develop structure-damaged joints, including joint space narrowing and osteophytes and the associated symptoms such as pain, stiffness, and ache. Structurally, OA is characterized by cartilage loss and modification of the structural and material properties of the subchondral bone.<sup>8</sup> Due to aging, obesity, and injury, cartilage loss in the joints gradually reduces the protective space between the bones, ultimately causing bone rubs against the bone. Earlier studies have suggested that knee cartilage in persons with knee OA tends to deteriorate more rapidly than those without the disease and have nearly double the rate of tibial cartilage loss among those with symptomatic knee OA than the healthy controls without knee pain.<sup>9,10</sup> These studies suggest a close relationship between cartilage loss and the progression and pain of knee OA.

Furthermore, a study by Hunter et al<sup>11</sup> suggested that meniscal damage was strongly associated with cartilage loss. The authors found that less meniscal coverage and reduced meniscal height were significantly associated with cartilage loss.<sup>11</sup> Consistent with this, another study reported that the morphology of the meniscal damage, especially root tear, was associated with the highest score of Western Ontario and McMaster Universities Osteoarthritis (WOMAC) pain score in the knees.<sup>12</sup> Again, these studies have illustrated meniscal damage as part of knee OA structural progression pathophysiology.

**ORCID iD of the author:**  
Z.D. 0000-0003-0809-5692.

**Cite this article as:** Dai Z. A literature review on plant-based foods and dietary quality in knee osteoarthritis. *Eur J Rheumatol* 2022;10.5152/eurjrheum.2022.21134 [Epub Ahead of Print].

<sup>1</sup> Charles Perkins Centre, School of Pharmacy, Faculty of Medicine and Health, The University of Sydney, Australia

<sup>2</sup> College of Medicine & Public Health Flinders University, Australia

Corresponding author:  
Zhaoli Dai  
E-mail: zhaoli.dai-keller@sydney.edu.au

Received: June 29, 2021  
Accepted: January 25, 2022

Available Online Date: April 22, 2022

Copyright © Author(s) - Available online at  
www.eurjrheumatol.org.

Content of this journal is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.



Osteophytes, another structural change of the joints, are often used to define the presence of OA radiographically. These bone spurs are most often present at the margins of the joint, as outgrowths of cartilage originally and subsequently undergo endochondral ossification.<sup>13</sup> Although osteophytes may not increase the risk of OA structural progression, they have been shown to be strongly associated with malalignment to the side of the osteophytes.<sup>13</sup> Other studies in the well-adjusted cross-sectional analyses also suggested that osteophyte size was associated with the presence<sup>14</sup> but not the severity of pain in knee OA.<sup>15</sup>

Besides structural damages found in OA, a growing body of evidence supports the notion that inflammation, both locally found in the joints as well as the low-grade systemic inflammation due to obesity or other metabolic diseases, may contribute to the structural progression of OA joints, pain, and symptoms.<sup>16</sup> As a form of local inflammation, synovitis is commonly found in the synovial membrane and it has been increasingly recognized as a pathophysiological mechanism of OA progression.<sup>17</sup> In support of this, evidence on analgesic therapies supports the anti-inflammatory agents for the benefits of targeting synovial inflammation to reduce knee pain.<sup>18</sup> Overall, this body of evidence suggests that synovial inflammation is associated with progressive joint failure,<sup>19,20</sup> incident OA,<sup>21,22</sup> and OA symptoms, particularly knee pain.<sup>23,24</sup>

### Main Points

- Plant-based food consumption such as fruit, vegetables, and legumes is likely to have a favorable outcome in knee osteoarthritis (OA), although the quality of the evidence is low.
- Studies on dietary quality measured by the Mediterranean diet, alternative healthy eating index, or the dietary inflammatory index also suggest possible protective associations with different clinical or pathophysiological indicators of OA.
- Potential mechanisms for associating these dietary components with knee osteoarthritis are likely contributing to the dietary effects on obesity, inflammation, and the gut microbiome. Human data in the microbiome–joint pain association are lacking but suggest that microbial composition could be a target to reduce inflammatory pain in OA.

Together, the interlinks of local (synovitis) and systemic inflammation with cartilage damage in knee OA joints demonstrate the potential of an anti-inflammatory intervention in managing the structural progression and knee pain related to knee OA.

Non-pharmacologic treatment options including physical activity, safe and monitored weight loss through dietary interventions, and physical therapy should be the first-line treatment options to manage knee OA.<sup>25</sup> The American College of Rheumatology guidelines strongly recommend weight loss to overweight or obese individuals with knee and hip OA.<sup>25</sup> This recommendation is based on a dose–response relationship between weight loss and improved symptoms and functions in patients with knee OA.<sup>26</sup> Later studies and reviews also have suggested that over 7.7%–10% of body weight loss has clinically meaningful outcomes in OA in the knees.<sup>27–29</sup>

Dietary quality plays a crucial role in overweight and obesity. The World Health Organization<sup>3</sup> and different countries' dietary guidelines recommend increased consumption of plant-based foods, including whole grains, fruit and vegetables, and legumes and nuts, as an effective way to control weight gain and reduce obesity. Furthermore, emerging evidence has suggested that plant-based foods or healthy diets may reduce chronic musculoskeletal pain,<sup>30</sup> including fibromyalgia<sup>31</sup> and rheumatoid arthritis.<sup>32</sup>

Earlier studies using data from different population-based longitudinal cohorts have shown the health benefits of fiber-rich foods in lowering the risk of symptomatic knee OA and knee pain severity.<sup>33</sup> This protective association is partly explained by the mechanism where fiber lowers bodyweight that might precede its effect on the symptoms due to knee OA. Furthermore, this protective association is potentially mediated by reducing inflammation, such as C-reactive protein (CRP).<sup>34</sup> Consistent with our findings, several healthy dietary patterns in the published systematic reviews, including a low inflammatory diet<sup>26</sup> and the Mediterranean diet,<sup>35</sup> have suggested a protective association with knee OA in observational studies or clinical trials. However, it should be noted that the heterogeneity of these studies included in these reviews is high.

To my knowledge, no review has synthesized the evidence on plant-based foods such as fruit, vegetables, nuts/legumes, whole grains, and diets that are rich in these components

regarding their roles in the prevention and treatment of knee OA. Therefore, the first part of the review summarizes these empirical findings. The second part explores the potential mechanisms that affect knee OA outcomes via body weight, inflammation, and the gut microbiome.

Briefly, a literature search using keywords, including osteoarthritis, plant-based, dietary quality, healthy diet, food, and knee osteoarthritis, were conducted in PubMed via Ovid to search articles published in 2015–2020. The inclusion criteria are human observational studies or clinical trials. The exposures or interventions are plant-based foods mentioned above or a healthy dietary pattern such as a Mediterranean diet or a diet based on a dietary quality index. The clinical outcomes must relate to structures or symptoms of knee OA. Studies on nutrients, such as dietary magnesium or vitamins including K and B complex, were excluded, as these nutritional sources vary and include both plant-based and animal foods. Studies focusing on nutrient supplements are also excluded. All included studies were written in English.

### Evidence on the Association of Diet and Knee OA

A total of 14 studies were included in this review. In addition, a narrative summary was provided based on study design, dietary exposures or interventions, and outcome measures. Table 1 describes the studies focusing on plant-based dietary components and their association with OA, and Table 2 summarizes studies focusing on dietary patterns with OA.

### Role of Plant-Based Foods in OA

In studies focusing on plant-based foods, there are 3 cross-sectional studies<sup>36–38</sup> and 5 randomized, controlled trials.<sup>39–43</sup> The exposures include soymilk (once a day vs. less than once a day), fruit or vegetables (times/day), and fruit and vegetable intakes in quantiles in the observational studies. The interventions include pomegranate juice (200 mL daily vs. usual lifestyle), broccoli (100 g/day vs. regular diets), freeze-dried strawberries (50 g/day vs. placebo), and freeze-dried blueberries (40 g/day vs. placebo). The clinical outcomes include cartilage structure parameters such as osteophytes (OST), joint space narrowing (JSN), cartilage defects, bone marrow lesions (BML), knee pain, WOMAC scores, and inflammatory markers such as IL-6, IL-1 $\beta$ , and matrix metalloproteinase (MMP-3). One study shows a higher level of synovial isothiocyanates in the broccoli treatment group than the control group as proof of concept of showing isothiocyanates

**Table 1.** Studies on the Health Benefits of Plant-Based Food Components in Knee or Hip Osteoarthritis

Study	Study Design	Populations	Food Components	Duration	Outcome Measures	Results	Favorable Outcomes or Not
<b>Observational studies</b>							
Li et al. 2016 (37)	Cross-sectional study	5764 people included	Soy milk intake	n/a	Prevalence of radiographic knee JSN and OST.	Significant associations found in those who had soy milk "≥once a day" vs. never: OR: 0.49, 95% CI: 0.26-0.92, P = .026 and those who had soy milk <once a day vs. never: OR: 0.79, 95% CI: 0.67-0.94, P = .009 and the P for trend was .001	Yes with OST but no association with JSN.
Wang et al. 2016 (38)	Cross-sectional study	214 participants without diagnosed hip OA	Fruit and vegetables	n/a	Femoral head BML and cartilage defects	Vegetable consumption (times/day) was associated with reduced prevalence of femoral head cartilage defects and BML. No associations for fruit consumption.	Yes for vegetable consumption with prevalence of femoral head cartilage defects and BML but no protective association with fruit consumption.
Han et al. 2017 (39)	Cross-sectional study	6588 people aged ≥50 years	Fruit and vegetable intake	n/a	Pain scores of knee decreased significantly with increasing fruit and vegetable intake by quartile.	Fourth versus the first quartile of vegetable and fruit consumption: decreased prevalence of severe knee pain (OR: 0.59, 95% CI: 0.48-0.73)	Fruit and vegetable intake decreased pain score.
<b>Clinical trials</b>							
Ghoochani et al. 2016 (40)	Randomized clinical trial	38 patients with knee OA	200 mL sugar- and additive-free pomegranate juice daily versus usual lifestyle (placebo)	6 weeks	Clinical signs on stiffness and functions, inflammation and antioxidant status.	Group differences were found for lower levels of MMP1, MMP13, and glutathione peroxidase in the treatment group versus the control group after the intervention. No group differences in WOMAC pain or other WOMAC scores. However, WOMAC pain score and stiffness were lower after the intervention in the treatment group.	Pomegranate juice may contain phytochemicals, such as anthocyanins and ellagic acid which may inhibit proteoglycan breakdown in cartilage, evident by decreasing MMPs; they may also exert anti-inflammatory and antioxidant properties.
Davidson et al. 2017 (41)	Randomized clinical trial (crossover)	40 patients with knee OA undergoing total knee replacement	7 days_ washout_ 100 g of broccoliid versus no such diet	14 days before TKR	Synovial isothiocyanates	Synovial ITC was detected in the high glycosylate diet but not in the low group; however, mean total ITC levels were 4.47-fold lower in the synovial fluid than in the plasma.	Bioactive constituent ITCs reach the synovial fluid at concentrations with biological impact on the articular joint tissues and alter the synovial fluid protein profile, showing that a dietary bioactive with chondroprotective properties can penetrate the knee in osteoarthritis.

(Continued)

**Table 1.** Studies on the Health Benefits of Plant-Based Food Components in Knee or Hip Osteoarthritis (Continued)

Study	Study Design	Populations	Food Components	Duration	Outcome Measures	Results	Favorable Outcomes or Not
Schell et al. 2017 (42)	Randomized clinical trial (crossover)	17 adults with knee OA	Freeze-dried strawberry beverage (50 g/day) or control beverage daily	12 weeks	IL-6, IL-1, MMP-3, pain score, and disability index	Interleukin (IL)-6, IL-1 $\beta$ , and MMP-3 were significantly decreased after consumption of strawberry beverage versus control (all $P < .05$ ). In addition, the strawberry group also significantly reduced constant, intermittent, and total pain as evaluated by the ICOAP questionnaire and the HAQ-DI scores as a disability index (all $P < .05$ ), compared with the control group.	Yes, with a reduction on IL-6, IL-1, MMP-3, pain score and disability index. No effects on CRP, lipid profiles.
Basu et al. 2018 (43)	Randomized clinical trial (crossover)	4 men and 13 women aged 57 $\pm$ 3 year	Dietary freeze-dried strawberries vs. control, 50 g/day (match for kcal and fiber)	12 weeks, washout 2 weeks	Obesity-related hormones, biomarkers of inflammation and lipid peroxidation	TNF- $\alpha$ and lipid peroxidation markers, hydroxy-2-nonenal (4-HNE) and conjugated dienes. decreased, but not body weight	Yes to TNF- $\alpha$ and lipid peroxidation markers but no effect on body weight.
Du et al. 2019 (44)	Randomized clinical trial	63 adults aged 45-79 with symptomatic knee OA	Freeze-dried whole blueberries (40 g/day) versus placebo on pain, gait performance, and inflammation in individuals with symptomatic knee OA	4 months	WOMAC pain, walking speed, and inflammation markers include IL-1, IL-6, IL-10, IL-13.	WOMAC total score and sub-groups, including pain, stiffness, and difficulty to perform daily activities, decreased significantly in the blueberry treatment group ( $P < .05$ ) but no difference from the control group. No effect was found on reducing inflammatory markers such as IL-1, IL-13, IL-6, and IL-10 or MCP-1 between time points in the same group or between groups.	No differences between groups.

OST, osteophytes; JSN, joint space narrowing; BML, bone marrow lesion; MMP, matrix metalloproteinase; OR, odds ratio; WOMAC, Western Ontario and McMaster Universities Osteoarthritis pain score; TKR, total knee replacement; ITC, iotrochyanates; CRP, C-reactive protein; MCP, monocyte chemoattractant protein-1; TNF, tumor necrosis factor; OA, osteoarthritis.

**Table 2.** Studies on Dietary Patterns in Association with Knee Osteoarthritis

Study	Study Design	Populations	Food Or		Duration	Outcome Measures	Results	Favorable Outcomes Or Not	
			Nutrients	Nutrients				Or	Not
Veronese et al. 2016 (45)	Cross-sectional studies	4470 participants aged 61 years on average	aMED		n/a	Quality of life by SF-12, WOMAC pain, and CES-D	Higher aMED was significantly associated with a higher SF-12 physical composite scale value (beta: 0.10; 95% CI: 0.05, 0.15; $P < .0001$ ), lower WOMAC scores (except for stiffness), and lower CES-D scores (beta: $-0.05$ ; 95% CI: $-0.09$ , $-0.01$ ; $P = .01$ )	Yes, to quality of life scores, WOMAC pain, and CES-D (mental health) score.	
Veronese et al. 2017 (46)	Cross-sectional studies	4358 community-dwelling participants (2527 females; mean age: 61.2 years) from OAI	aMED		n/a	Prevalence of knee OA	Participants with the highest aMED had a significant reduction in the presence of knee OA (OR, 0.83; 95% CIs: 0.69-0.99, $P = .04$ ). In addition, higher use of cereals was associated with lower odds of having knee OA (OR: 0.76; 95% CI: 0.60-0.98; $P = .03$ ).	Yes, to lower the prevalence of knee OA.	
Mears et al. 2018 (47)	Cross-sectional studies	122 older overweight and obese African American females with self-reported OA	Alternative Healthy Eating Index 2010 (AHEI-2010)		n/a	WOMAC, IL-6	Interleukin-6 was negatively correlated with AHEI-2010	Yes, to IL-6	
Veronese 2018 (48)	Cross-sectional studies	783 participants with an MRI assessment	MED		n/a	Knee cartilage morphology	Per standard deviation in the MED corresponded to a significant increase in the central medial femoral cartilage volume (beta = 0.12; 95% CI: 0.09-0.15), in the mean central medial femoral cartilage thickness (beta = 0.13; 95% CI: 0.01-0.17), in the cartilage thickness of the mean central medial tibiofemoral compartment (beta = 0.12; 95% CI: 0.09-0.15), and in the cartilage volume of the medial tibiofemoral compartment (beta = 0.09; 95% CI: 0.06-0.12).	Yes, to increase cartilage volume and cartilage thickness.	
Veronese et al. 2019 (49)	Cross-sectional studies	4358 participants	DII		n/a	Radiographic, symptomatic knee OA	Participants with the highest versus lowest quartile DII, had a significantly higher probability of experiencing radiographic symptomatic knee osteoarthritis (OR: 1.40; 95% CI: 1.14-1.72; $P = .002$ ).	A negative effect of an inflammatory diet to knee OA	
<b>Clinical trials</b>									
Dyer et al. 2017 (50)	Randomized trial	99 volunteers with OA (aged 31-90 years) (med diet: n = 50 versus normal diet n = 49)	Mediterranean diet		16 weeks	AIMS2 and inflammatory cytokines	There was no difference in AIMS2 and most inflammatory markers between the groups, except IL-1 alpha and sCOMP decreased in the diet group, but a significant improvement in the hip rotation in the diet group versus the control group.	Yes, to IL-1 and sCOMP and hip rotation. No effects on most inflammatory markers or arthritis impact measurement scale.	

CES-D, Center for Epidemiological Studies Depression; aMED, alternative Mediterranean diet; WOMAC, Western Ontario and McMaster Universities Osteoarthritis pain score; OR, odds ratio; OA, osteoarthritis; MRI, magnetic resonance imaging; DII, Dietary Inflammatory Index; IL, interleukin; sCOMP, serum cartilage oligomeric matrix protein; AIMS2, Arthritis Impact Measurement Scale; SF-12, Short Form Survey 12.

could penetrate to the synovial fluid. However, this study did not mention its effect on OA outcomes.<sup>40</sup>

Half of the studies have suggested favorable outcomes on symptoms (knee pain, stiffness, or symptomatic knee OA) or structure (JSN, OST, and radiographic knee OA). To be noted, the sample size of the intervention studies is generally small, ranging from 17 to 63 individuals with a trial period of 14 days to 4 months. In addition, although significant effects were reported on reduced pain or other symptoms<sup>39,43</sup> or lower levels of IL-1, IL-6, TNF, and MMP-3 markers,<sup>41,42</sup> no consistent findings were reported across the studies.

#### Role of Dietary Patterns/Diet Quality in OA

In studies focusing on dietary patterns and their associations with or effects on knee OA-related outcomes, there are 5 cross-sectional studies<sup>44-48</sup> and 1 RCT<sup>49</sup> (Table 2). The majority of the studies examined the Mediterranean diet (n=5). Two studies assessed diet quality, one on the Alternative Healthy Eating Index 2010 and the other examined the dietary inflammatory index (DII). The outcomes include the prevalence of OA, quality of life using the 12-Item Short Form Survey (SF-12), WOMAC pain, Center for Epidemiological Studies Depression (CES-D), levels of inflammatory cytokines such as IL-1, serum cartilage oligomeric matrix protein (sCOMP), and structural outcomes on cartilage volume/thickness. All studies found significant associations for all or partial indicators of OA. The only RCT among 99 participants suggests that those who consumed the Mediterranean diet had a reduction in IL-1 and sCOMP and a significant improvement in hip rotation than those on the control diet after a 16 week of intervention.<sup>49</sup>

#### Clinical Relevance of the Included Intervention Studies

Although the results of the 6 included intervention studies have indicated certain health benefits of the dietary patterns to manage OA, the statistically significant differences between the control and the treatment groups were primarily found based on the change in the inflammatory or oxidative markers.<sup>39,41,42,49</sup> However, no statistical significance was found in group differences for WOMAC pain,<sup>39</sup> bodyweight,<sup>42</sup> or Arthritis Impact Measurement Scale.<sup>49</sup> Furthermore, in some studies, the authors only presented the changes between different time points in the treatment group instead of comparing the group differences.<sup>43</sup> Interestingly, the analysis by Davidson et al<sup>40</sup> has detected a higher concentration of isothiocyanates in the group of people who consumed broccoli,

suggesting specific dietary components may penetrate the articular joint tissue to alter the protein profile in the synovial fluid.

Because of the small sample size and a short intervention period, it remains to be determined whether these plant-based components have clinically meaningful benefits in OA parameters, such as cartilage degradation, joint space narrowing, pain, and stiffness. However, as pomegranate juice and berries contain polyphenols such as anthocyanins and ellagic acids that were shown to have anti-inflammatory and antioxidant functions, they were likely to inhibit various interleukine cytokines, serum cartilage oligomeric matrix protein (a marker of cartilage degradation), peroxidation in the chondrocytes, synoviocytes, macrophages, and osteoblast to have a downstream effect in reducing cartilage degradation, joint space narrowing, pain, and stiffness in OA.<sup>50</sup>

#### Comparison with Other Reviews on Food/Diets on OA:

This review is unique because it focuses on plant-based foods and dietary patterns and their relationship with OA onset, prevalence, related pain, and other pathophysiological parameters. We did not include nutritional supplements in this review for the following reasons. First, the supplements regulations are usually lax, and the quality of dietary supplements is generally low or unknown. Second, the evidence on the health benefits of supplements to maintain well-being remains controversial in different health conditions, including knee OA.<sup>51</sup> Third, a previous review on the effects of dietary supplements on OA management has pointed to low quality of evidence. In the review led by Liu and colleagues<sup>52</sup>, the authors reported that several unusual supplements such as *Boswellia serrata* extract and pycnogenol had significant effects on pain reduction in hand, knee, or hip OA. However, the most widely used supplements such as glucosamine, chondroitin, vitamin D, or vitamin E did not provide clinically meaningful long-term effects to improve pain symptoms.<sup>52</sup>

Another review on polyphenols primarily conducted in animal studies has provided evidence of their potential benefits on reducing levels of IL-1, IL-6, TNF, and MMP.<sup>53</sup> Consistent with this review, a meta-analysis on mice induced with OA by surgical procedures has suggested adverse outcomes on OA in the mice fed a high-fat diet, including worsened progression of OA and increased levels of TNF- $\alpha$ , IL-1 $\beta$ , and leptin.<sup>54</sup> However, there is only one systematic review on the Mediterranean diet,<sup>35</sup> which only includes two cross-sectional

studies<sup>44,45</sup> and one RCT.<sup>49</sup> Due to the high heterogeneity of the studies, no meta-analysis was conducted. Another systematic review focused on the inflammatory aspect of diets, including the Mediterranean diet and diet supplemented with blueberry or strawberry powders on knee pain related to rheumatoid arthritis and OA.<sup>26</sup> In this review, the meta-analysis has suggested statistically significant differences in weight change and inflammatory biomarkers, such as in CRP, IL-6, and IL-1 $\beta$ ; but no significant effects on physical function or pain score were detected. Again, heterogeneity across the studies was high (>70%). Overall, the quality of the evidence was rated low, and each meta-analysis included only 3 studies.<sup>26</sup>

#### Potential Mechanisms for Plant-Based Foods and Diet Quality on Knee OA

##### Overweight and Obesity

Overweight and obesity are the causative risk factors for knee OA. The weight-induced mechanical loading further exacerbates the age-related “wear and tear” process of structural changes in joint cartilage. Knee OA shares similar metabolic risk factors such as body weight and inflammation with several cardio-metabolic conditions, including obesity, type 2 diabetes, and cardiovascular diseases.<sup>55</sup> The lifetime risk of diagnosed knee OA was estimated as 20% for obese people compared to 11% for non-obese people, with the highest risk (24%) found in obese females.<sup>56</sup>

The effect of a low-calorie diet, including high consumption of vegetables, on knee OA, was well illustrated in an 18-month intervention, which found that a very low-calorie diet (500-700 kcal/day, high in vegetables) reduced knee joint compressive force in the diet arm or the diet+exercise arm and comparable lower levels of IL-6 than the exercise arm.<sup>57</sup> This evidence suggests that a diet rich in plant-based foods or food components with weight control effects is likely to relieve pain symptoms related to OA. Furthermore, a network meta-analysis has indicated that the most effective interventions to alleviate arthritis pain were bariatric surgery, low-calorie diet and exercise, and intensive weight loss and exercise.<sup>58</sup> For most people, consuming a diet rich in plant-based foods may offer a safe, economical, and sustainable option to prevent and manage OA if that is feasible.

##### Inflammation

Evidence on the health benefits of plant-based foods, such as fruit, vegetables, whole grains, nuts, and legumes lowering systemic inflammatory cytokines, has been studied

extensively.<sup>59,60</sup> For different healthy diets, such as the Mediterranean diet [emphasizing on fruits, vegetables, olive oil, cereals, nuts and legumes, and limited consumption of red meat, processed meat and full-fat dairy<sup>61</sup>], the alternative healthy eating index 2010 [focusing on fruits, vegetables, whole grains, nuts and legumes, and long-chain unsaturated fatty acids, and minimized red meat and processed meat consumption<sup>62</sup>], and the DII [including several herbs and spices, dietary fiber, vitamins, fatty acids, polyphenols such as genistein, daidzein, and anthocyanin, carbohydrates, and alcohol; with the higher the score, the more pro-inflammatory the diet is<sup>63</sup>], several systematic reviews have demonstrated their health benefits in mortality,<sup>64,65</sup> cardiovascular diseases, and type 2 diabetes.<sup>65</sup> In addition, higher DII is related to a higher risk of developing central obesity<sup>66</sup> and mortality.<sup>67</sup> These specific dietary patterns (for DII, a diet with a low inflammatory score) may affect inflammation through multiple pathways, such as DNA methylation to modulate gene expression,<sup>68</sup> lower oxidative stress,<sup>69</sup> improve autophagy,<sup>70</sup> and regulate helper T cells in the immune response<sup>71</sup> that lead to a downstream effect to reduce inflammation.

## Microbiome

### Diet and Microbiome

Another potential mechanism is the effect of plant-based diets on the microbiome's composition. The relationship between diet and the microbiome has been established with convincing evidence.<sup>72-75</sup> Several reviews have also reported that plant-based diets, especially those containing higher contents of dietary fiber and polyphenols, may promote a more diverse and stable microbial system. The health benefits were seen as higher counts of microbes with anti-pathogenic and anti-inflammatory effects, including the lactic acid bacteria (e.g., *Ruminococcus*, *Eubacterium rectale*, and *Roseburia*), *Bifidobacterium* and *Lactobacillus*, and reduced counts of *Clostridium* and *Enterococcus* genera. Furthermore, the metabolites of dietary fiber after fermentation, such as short-chain fatty acids (acetate, propionate, and butyrate), can also improve immunity against pathogens and further regulate critical function of the intestine.<sup>75</sup>

In human studies, a cohort has shown a lower relative abundance of the phylum *Actinobacteria* (a group of gram-positive bacteria) in older adults of different ethnic backgrounds after they consumed a high-quality diet based on the Healthy Eating Index, Alternative Healthy Eating Index-2010,

the Mediterranean Diet, and the Dietary Approaches to Stop Hypertension.<sup>76</sup> For example, of the 104 bacterial genera tested, 21 (primarily from the phylum *Firmicutes*) were positively associated with at least one dietary index.<sup>76</sup> This evidence suggests that diet quality may link to fecal microbial alpha and beta diversity and different genera. In addition, the increased production of short-chain fatty acids may suppress the activation of effector T cells and abrogate the manifestation of undesirable inflammatory responses.<sup>77</sup> However, whether this is the case in knee OA or OA in general, remains tested.

### Microbiome and OA

The gut microbiome concerning health and disease development is still a relatively new research area, particularly OA. One piece of evidence has shown a correlation between dysbiosis of microbiota and the pathogenesis of OA. It should be noted that the majority of the studies were conducted in animal studies. One such review has suggested favorable effects on reduced joint pain and inflammatory markers after feeding either prebiotic or probiotics or *Lactobacillus* to OA-induced mice or rats.<sup>78,79</sup> However, human data in this aspect are limited. A Rotterdam study including more than 1000 participants with hip and knee OA has found a positive relationship between a higher WOMAC score (more pain) and the abundance of the pro-inflammatory *Streptococcus* taxa.<sup>80</sup> A follow-up of this study, which includes the Rotterdam cohort and the validation cohort of older Canadians, has further confirmed the independent association between greater relative and absolute *Streptococcus* spp. abundance and higher knee pain and local joint inflammation.<sup>81</sup> Additionally, a Chinese cohort of 125 patients with rheumatoid arthritis and 58 patients with OA found abundant diversity of bacterial nucleic acids correlated with the disease condition and that patients with OA had more abundant pro-inflammatory *Haemophilus parainfluenzae*, *Bacteroides fragilis*, *Porphyromonas*, and *Streptococcus* in their synovial tissues or synovial fluid.<sup>82</sup> In summary, these pioneer human observational studies have established a positive relationship between the gut microbiome and OA pathophysiology, suggesting that the microbiome could be a potential therapeutic target for interventions to manage OA pain.

## Conclusion

This literature review summarizes the most relevant and updated studies on the role of plant-based food and diet quality in knee

OA in observational studies and clinical trials. Although the evidence has shown a protective association with pain, cartilage structural parameters, and inflammatory biomarkers related to OA, the quality of this body of evidence is generally low. Furthermore, as there are considerable variations in the results across the studies, high-quality longitudinal cohort studies and well-designed clinical trials are needed to ascertain the causal effects of plant-based food on clinically meaningful outcomes in OA. In addition, a limited number of observational studies have indicated a potential link between the microbiome and OA. Finally, given the close relationships of plant-based food on obesity, inflammation, and microbiome, evidence on the role of these foods and diet concerning knee OA pain through the changes in the microbial composition can provide more certainty to explore this therapeutic target.

**Peer-Review:** Externally peer-reviewed.

**Acknowledgments:** The author would like to acknowledge Richard L. Cohen for proofreading the manuscript. Richard is a longtime health care journalist and previously was the editor of Healthcare Marketing Report and Healthcare Call Center Times.

**Declaration of Interests:** The author has no conflict of interest to declare.

**Funding:** The author declared that this study has received no financial support.

## References

1. GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1789-1858. [\[CrossRef\]](#)
2. Kloppenburg M, Berenbaum F. Osteoarthritis year in review 2019: epidemiology and therapy. *Osteoarthr Cartil*. 2020;28(3):242-248. [\[CrossRef\]](#)
3. The World Health Organization. Obesity and overweight; 2021. Available at: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>.
4. Hunter DJ, March L, Chew M. Osteoarthritis in 2020 and beyond: a lancet commission. *Lancet*. 2020;396(10264):1711-1712. [\[CrossRef\]](#)
5. Berenbaum F, Wallace IJ, Lieberman DE, Felson DT. Modern-day environmental factors in the pathogenesis of osteoarthritis. *Nat Rev Rheumatol*. 2018;14(11):674-681. [\[CrossRef\]](#)
6. Cicuttini FM, Wluka AE. Not just loading and age: the dynamics of osteoarthritis, obesity and inflammation. *Med J Aust*. 2016;204(2):47. [\[CrossRef\]](#)

7. Wang X, Hunter D, Xu J, Ding C. Metabolic triggered inflammation in osteoarthritis. *Osteoarthritis Cartil.* 2015;23(1):22-30. [\[CrossRef\]](#)
8. Cox LG, van Rietbergen B, van Donkelaar CC, Ito K. Bone structural changes in osteoarthritis as a result of mechanoregulated bone adaptation: a modeling approach. *Osteoarthritis Cartil.* 2011;19(6):676-682. [\[CrossRef\]](#)
9. Hanna F, Ebeling PR, Wang Y, et al. Factors influencing longitudinal change in knee cartilage volume measured from magnetic resonance imaging in healthy men. *Ann Rheum Dis.* 2005;64(7):1038-1042. [\[CrossRef\]](#)
10. Wluka AE, Wolfe R, Davis SR, Stuckey S, Cicuttini FM. Tibial cartilage volume change in healthy postmenopausal women: a longitudinal study. *Ann Rheum Dis.* 2004;63(4):444-449. [\[CrossRef\]](#)
11. Hunter DJ, Zhang YQ, Niu JB, et al. The association of meniscal pathologic changes with cartilage loss in symptomatic knee osteoarthritis. *Arthritis Rheum.* 2006;54(3):795-801. [\[CrossRef\]](#)
12. MacFarlane LA, Yang H, Collins JE, et al. Associations among meniscal damage, meniscal symptoms and knee pain severity. *Osteoarthritis Cartil.* 2017;25(6):850-857. [\[CrossRef\]](#)
13. Felson DT, Gale DR, Elon Gale M, et al. Osteophytes and progression of knee osteoarthritis. *Rheumatol (Oxf Engl).* 2005;44(1):100-104. [\[CrossRef\]](#)
14. Kornaat PR, Bloem JL, Ceulemans RY, et al. Osteoarthritis of the knee: association between clinical features and MR imaging findings. *Radiology.* 2006;239(3):811-817. [\[CrossRef\]](#)
15. Sengupta M, Zhang YQ, Niu JB, et al. High signal in knee osteophytes is not associated with knee pain. *Osteoarthritis Cartil.* 2006;14(5):413-417. [\[CrossRef\]](#)
16. Scanzello CR, Loeser RF. Editorial: inflammatory activity in symptomatic knee osteoarthritis: not all inflammation is local. *Arthritis Rheumatol.* 2015;67(11):2797-2800. [\[CrossRef\]](#)
17. Sokolove J, Lepus CM. Role of inflammation in the pathogenesis of osteoarthritis: latest findings and interpretations. *Ther Adv Musculoskelet Dis.* 2013;5(2):77-94. [\[CrossRef\]](#)
18. van Laar M, Pergolizzi JV, Jr, Mellinshoff HU, et al. Pain treatment in arthritis-related pain: beyond NSAIDs. *Open Rheumatol J.* 2012;6:320-330. [\[CrossRef\]](#)
19. Damman W, Liu R, Bloem JL, Rosendaal FR, Rejniers M, Kloppenburg M. Bone marrow lesions and synovitis on MRI associate with radiographic progression after 2 years in hand osteoarthritis. *Ann Rheum Dis.* 2017;76(1):214-217. [\[CrossRef\]](#)
20. Haugen IK, Slatkowsky-Christensen B, Bøyesen P, Sesseng S, van der Heijde D, Kvien TK. MRI findings predict radiographic progression and development of erosions in hand osteoarthritis. *Ann Rheum Dis.* 2016;75(1):117-123. [\[CrossRef\]](#)
21. Atukorala I, Kwok CK, Guerazzi A, et al. Synovitis in knee osteoarthritis: a precursor of disease? *Ann Rheum Dis.* 2016;75(2):390-395. [\[CrossRef\]](#)
22. Roemer FW, Kwok CK, Hannon MJ, et al. What comes first? Multitissue involvement leading to radiographic osteoarthritis: magnetic resonance imaging-based trajectory analysis over four years in the osteoarthritis initiative. *Arthritis Rheumatol.* 2015;67(8):2085-2096. [\[CrossRef\]](#)
23. Hill CL, Hunter DJ, Niu J, et al. Synovitis detected on magnetic resonance imaging and its relation to pain and cartilage loss in knee osteoarthritis. *Ann Rheum Dis.* 2007;66(12):1599-1603. [\[CrossRef\]](#)
24. Zhang Y, Nevitt M, Niu J, et al. Fluctuation of knee pain and changes in bone marrow lesions, effusions, and synovitis on magnetic resonance imaging. *Arthritis Rheum.* 2011;63(3):691-699. [\[CrossRef\]](#)
25. Kolasinski SL, Neogi T, Hochberg MC, et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the management of osteoarthritis of the hand, hip, and knee. *Arthritis Care Res (Hoboken)* 2020;72(2):149-162. [\[CrossRef\]](#)
26. Genel F, Kale M, Pavlovic N, Flood VM, Naylor JM, Adie S. Health effects of a low-inflammatory diet in adults with arthritis: a systematic review and meta-analysis. *J Nutr Sci.* 2020;9:e37. [\[CrossRef\]](#)
27. Messier SP, Resnik AE, Beavers DP, et al. Intentional weight loss in overweight and obese patients With knee osteoarthritis: is more better? *Arthritis Care Res (Hoboken).* 2018;70(11):1569-1575. [\[CrossRef\]](#)
28. Jin X, Gibson AA, Gale J, et al. Does weight loss reduce the incidence of total knee and hip replacement for osteoarthritis? A prospective cohort study among middle-aged and older adults with overweight or obesity. *Int J Obes (Lond).* 2021;45(8):1696-1704. [\[CrossRef\]](#)
29. Atukorala I, Makovey J, Lawler L, Messier SP, Bennell K, Hunter DJ. Is there a dose-response relationship between weight loss and symptom improvement in persons with knee osteoarthritis? *Arthritis Care Res (Hoboken).* 2016;68(8):1106-1114. [\[CrossRef\]](#)
30. Elma Ö, Yilmaz ST, Delihs T, et al. Do nutritional factors interact with chronic musculoskeletal pain? A systematic review. *J Clin Med.* 2020;9(3). [\[CrossRef\]](#)
31. Towery P, Guffey JS, Doerflin C, Stroup K, Saucedo S, Taylor J. Chronic musculoskeletal pain and function improve with a plant-based diet. *Complement Ther. Med.* 2018;40:64-69. [\[CrossRef\]](#)
32. McDougall J, Bruce B, Spiller G, Westerdahl J, McDougall M. Effects of a very low-fat, vegan diet in subjects with rheumatoid arthritis. *J Altern Complement Med.* 2002;8(1):71-75. [\[CrossRef\]](#)
33. Dai Z, Lu N, Niu J, Felson DT, Zhang Y. Dietary fiber intake in relation to knee pain trajectory. *Arthritis Care Res (Hoboken).* 2017;69(9):1331-1339. [\[CrossRef\]](#)
34. Dai Z, Jafarzadeh SR, Niu J, et al. Body mass index mediates the association between dietary fiber and symptomatic knee osteoarthritis in the osteoarthritis initiative and the Framingham osteoarthritis study. *J Nutr.* 2018;148(12):1961-1967. [\[CrossRef\]](#)
35. Morales-Ivorra I, Romera-Baures M, Roman-Viñas B, Serra-Majem L. Osteoarthritis and the Mediterranean diet: a systematic review. *Nutrients.* 2018;10(8):7. [\[CrossRef\]](#)
36. Li H, Zeng C, Wei J, et al. Relationship between soy milk intake and radiographic knee joint space narrowing and osteophytes. *Rheumatol Int.* 2016;36(9):1215-1222. [\[CrossRef\]](#)
37. Wang Y, Smith S, Teichtahl AJ, et al. Association Between dietary intake of antioxidants and prevalence of femoral head cartilage defects and bone marrow lesions in community-based adults. *J Rheumatol.* 2016;43(10):1885-1890. [\[CrossRef\]](#)
38. Han HS, Chang CB, Lee DC, Lee JY. Relationship between total fruit and vegetable intake and self-reported knee pain in older adults. *J Nutr Health Aging.* 2017;21(7):750-758. [\[CrossRef\]](#)
39. Ghoochani N, Karandish M, Mowla K, Haghhighizadeh MH, Jalali MT. The effect of pomegranate juice on clinical signs, matrix metalloproteinases and antioxidant status in patients with knee osteoarthritis. *J Sci Food Agric.* 2016;96(13):4377-4381. [\[CrossRef\]](#)
40. Davidson R, Gardner S, Jupp O, et al. Isothiocyanates are detected in human synovial fluid following broccoli consumption and can affect the tissues of the knee joint. *Sci Rep.* 2017;7(1):3398. [\[CrossRef\]](#)
41. Schell J, Scofield RH, Barrett JR, et al. Strawberries improve pain and inflammation in obese adults with radiographic evidence of knee osteoarthritis. *Nutrients.* 2017;9(9):28. [\[CrossRef\]](#)
42. Basu A, Kurien BT, Tran H, et al. Strawberries decrease circulating levels of tumor necrosis factor and lipid peroxides in obese adults with knee osteoarthritis. *Food Funct.* 2018;9(12):6218-6226. [\[CrossRef\]](#)
43. Du C, Smith A, Avalos M, et al. Blueberries improve pain, gait performance, and inflammation in individuals with symptomatic knee osteoarthritis. *Nutrients.* 2019;11(2):29. [\[CrossRef\]](#)
44. Veronese N, Stubbs B, Noale M, Solmi M, Luchini C, Maggi S. Adherence to the Mediterranean diet is associated with better quality of life: data from the osteoarthritis Initiative. *Am J Clin Nutr.* 2016;104(5):1403-1409. [\[CrossRef\]](#)
45. Veronese N, Stubbs B, Noale M, et al. Adherence to a Mediterranean diet is associated with lower prevalence of osteoarthritis: data from the osteoarthritis initiative. *Clin Nutr.* 2017;36(6):1609-1614. [\[CrossRef\]](#)
46. Mears M, Tussing-Humphreys L, Cerwinske L, et al. Associations between alternate healthy eating Index-2010, body composition, osteoarthritis severity, and interleukin-6 in older overweight and obese African American females with self-reported osteoarthritis. *Nutrients.* 2018;11(1):22. [\[CrossRef\]](#)
47. Veronese N, La Tegola L, Crepaldi G, Maggi S, Rogoli D, Guglielmi G. The association between



- the Mediterranean diet and magnetic resonance parameters for knee osteoarthritis: data from the osteoarthritis Initiative. *Clin Rheumatol*. 2018;37(8):2187-2193. [\[CrossRef\]](#)
48. Veronese N, Shivappa N, Stubbs B, et al. The relationship between the dietary inflammatory index and prevalence of radiographic symptomatic osteoarthritis: data from the osteoarthritis Initiative. *Eur J Nutr*. 2019;58(1):253-260. [\[CrossRef\]](#)
  49. Dyer J, Davison G, Marcora SM, Mauger AR. Effect of a Mediterranean type diet on inflammatory and cartilage degradation biomarkers in patients with osteoarthritis. *J Nutr Health Aging*. 2017;21(5):562-566. [\[CrossRef\]](#)
  50. Ansari MY, Ahmad N, Haqqi TM. Oxidative stress and inflammation in osteoarthritis pathogenesis: role of polyphenols. *Biomed Pharmacother*. 2020;129:110452. [\[CrossRef\]](#)
  51. Binns CW, Lee MK, Lee AH. Problems and prospects: public health regulation of dietary supplements. *Annu Rev Public Health*. 2018;39(1):403-420. [\[CrossRef\]](#)
  52. Liu X, Machado GC, Eyles JP, Ravi V, Hunter DJ. Dietary supplements for treating osteoarthritis: a systematic review and meta-analysis. *Br J Sports Med*. 2018;52(3):167-175. [\[CrossRef\]](#)
  53. Valsamidou E, Gioxari A, Amerikanou C, Zoumpoulakis P, Skarpas G, Kaliora AC. Dietary interventions with polyphenols in osteoarthritis: a systematic review directed from the preclinical data to randomized clinical studies. *Nutrients*. 2021;13(5). [\[CrossRef\]](#)
  54. Sansone V, Applefield RC, De Luca P, et al. Does a high-fat diet affect the development and progression of osteoarthritis in mice?: a systematic review. *Bone Joint Res*. 2019;8(12):582-592. [\[CrossRef\]](#)
  55. Dickson BM, Roelofs AJ, Rochford JJ, Wilson HM, De Bari C. The burden of metabolic syndrome on osteoarthritic joints. *Arthritis Res Ther*. 2019;21(1):289. [\[CrossRef\]](#)
  56. Losina E, Weinstein AM, Reichmann WM, et al. Lifetime risk and age at diagnosis of symptomatic knee osteoarthritis in the US. *Arthritis Care Res (Hoboken)*. 2013;65(5):703-711. [\[CrossRef\]](#)
  57. Messier SP, Mihalko SL, Legault C, et al. Effects of intensive diet and exercise on knee joint loads, inflammation, and clinical outcomes among overweight and obese adults with knee osteoarthritis: the IDEA randomized clinical trial. *JAMA*. 2013;310(12):1263-1273. [\[CrossRef\]](#)
  58. Panunzi S, Maltese S, De Gaetano A, Capristo E, Bornstein SR, Mingrone G. Comparative efficacy of different weight loss treatments on knee osteoarthritis: a network meta-analysis. *Obes Rev*. 2021;22(8):e13230. [\[CrossRef\]](#)
  59. Watzl B. Anti-inflammatory effects of plant-based foods and of their constituents. *Int J Vitam Nutr Res*. 2008;78(6):293-298. [\[CrossRef\]](#)
  60. Tsigalou C, Konstantinidis T, Paraschaki A, Stavropoulou E, Voidarou C, Bezirtzoglou E. Mediterranean diet as a tool to combat inflammation and chronic diseases. *An Overview Biomed*. 2020;8(7).
  61. Trichopoulou A, Lagiou P. Healthy traditional Mediterranean diet: an expression of culture, history, and lifestyle. *Nutr Rev*. 1997;55(11 Pt 1):383-389. [\[CrossRef\]](#)
  62. Chiuve SE, Fung TT, Rimm EB, et al. Alternative dietary indices both strongly predict risk of chronic disease. *J Nutr*. 2012;142(6):1009-1018. [\[CrossRef\]](#)
  63. Cavicchia PP, Steck SE, Hurley TG, et al. A new dietary inflammatory index predicts interval changes in serum high-sensitivity C-reactive protein. *J Nutr*. 2009;139(12):2365-2372. [\[CrossRef\]](#)
  64. Soltani S, Jayedi A, Shab-Bidar S, Becerra-Tomás N, Salas-Salvadó J. Adherence to the Mediterranean diet in relation to all-cause mortality: a systematic review and dose-response meta-analysis of prospective cohort studies. *Adv Nutr*. 2019;10(6):1029-1039. [\[CrossRef\]](#)
  65. Morze J, Danielewicz A, Hoffmann G, Schwing-shackl L. Diet quality as assessed by the healthy eating index, alternate healthy eating index, dietary approaches to stop hypertension score, and health outcomes: a second update of a systematic review and meta-analysis of cohort studies. *J Acad Nutr Diet*. 2020;120(12):1998-2031.e15. [\[CrossRef\]](#)
  66. Farhangi MA, Vajdi M. The association between dietary inflammatory index and risk of central obesity in adults: an updated systematic review and meta-analysis. *Int J Vitam Nutr Res*. 2020;90(5-6):535-552. [\[CrossRef\]](#)
  67. Zahedi H, Djalalinia S, Asayesh H, et al. A higher dietary inflammatory index score is associated with a higher risk of incidence and mortality of cancer: a comprehensive systematic review and meta-analysis. *Int J Prev Med*. 2020;11:15. [\[CrossRef\]](#)
  68. Jones PA, Takai D. The role of DNA methylation in mammalian epigenetics. *Science*. 2001;293(5532):1068-1070. [\[CrossRef\]](#)
  69. Hussain T, Tan B, Yin Y, Blachier F, Tossou MCB, Rahu N. Oxidative stress and inflammation: what polyphenols can do for us? *Oxid Med Cell Longev*. 2016;2016:1-9. [\[CrossRef\]](#)
  70. Levine B, Mizushima N, Virgin HW. Autophagy in immunity and inflammation. *Nature*. 2011;469(7330):323-335. [\[CrossRef\]](#)
  71. Skapenko A, Leipe J, Lipsky PE, Schulze-Koops H. The role of the T cell in autoimmune inflammation. *Arthritis Res Ther*. 2005;7(suppl 2):S4-S14. [\[CrossRef\]](#)
  72. David LA, Maurice CF, Carmody RN, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature*. 2014;505(7484):559-563. [\[CrossRef\]](#)
  73. Kolodziejczyk AA, Zheng D, Elinav E. Diet-microbiota interactions and personalized nutrition. *Nat Rev Microbiol*. 2019;17(12):742-753. [\[CrossRef\]](#)
  74. Leeming ER, Louca P, Gibson R, Menni C, Spector TD, Le Roy CI. The complexities of the diet-microbiome relationship: advances and perspectives. *Genome Med*. 2021;13(1):10. [\[CrossRef\]](#)
  75. Tomova A, Bukovsky I, Rembert E, et al. The effects of vegetarian and vegan diets on gut microbiota. *Front Nutr*. 2019;6:47. [\[CrossRef\]](#)
  76. Maskarinec G, Hullar MAJ, Monroe KR, et al. Fecal microbial diversity and structure are associated with diet quality in the multiethnic cohort adiposity phenotype study. *J Nutr*. 2019;149(9):1575-1584. [\[CrossRef\]](#)
  77. Clemente JC, Manasson J, Scher JU. The role of the gut microbiome in systemic inflammatory disease. *BMJ*. 2018;360:j5145. [\[CrossRef\]](#)
  78. Biver E, Berenbaum F, Valdes AM, et al. Gut microbiota and osteoarthritis management: an expert consensus of the European society for clinical and economic aspects of osteoporosis, osteoarthritis and musculoskeletal diseases (ESCEO). *Ageing Res Rev*. 2019;55:100946. [\[CrossRef\]](#)
  79. Guo R, Chen LH, Xing C, Liu T. Pain regulation by gut microbiota: molecular mechanisms and therapeutic potential. *Br J Anaesth*. 2019;123(5):637-654. [\[CrossRef\]](#)
  80. Ulici V, Kelley KL, Azcarate-Peril MA, et al. Osteoarthritis induced by destabilization of the medial meniscus is reduced in germ-free mice. *Osteoarthritis Cartil*. 2018;26(8):1098-1109. [\[CrossRef\]](#)
  81. Boer CG, Radjabzadeh D, Medina-Gomez C, et al. Intestinal microbiome composition and its relation to joint pain and inflammation. *Nat Commun*. 2019;10(1):4881. [\[CrossRef\]](#)
  82. Zhao Y, Chen B, Li S, et al. Detection and characterization of bacterial nucleic acids in culture-negative synovial tissue and fluid samples from rheumatoid arthritis or osteoarthritis patients. *Sci Rep*. 2018;8(1):14305.